

Summary of Fire Risk Classes

Low (Green): The combination of the local fuel hazard, weather influences, topography, proximity to the community, fuel position in relation to fire spread patterns, and known local wildfire threat factors make it a lower potential for threatening a community. These stands will support surface fires, single tree or small groups of conifer trees could torch/ candle in extreme fire weather conditions. Fuel type spot potential is very low, low risk to any values at risk.

Moderate (Yellow): The combination of the local fuel hazard, weather influences, topography, proximity to the community, fuel position in relation to fire spread patterns and known local wildfire threat factors make it possible that a wildfire in this area would threaten the community. Areas of matted grass, slash, conifer plantations, mature conifer stands with very high crown base height, and deciduous stands with 26 to 49% conifers. These stands will support surface fires, single tree or small groups of conifer trees could torch/ candle. Rates of spread would average between 2-5 meters/ minute. Forest stands would have potential to impact values in extreme weather conditions. Fuel type spot potential is unlikely to impact values at a long distance (<400m).

High (Orange): The combination of the local fuel hazard, weather influences, topography, proximity to the community, fuel position in relation to fire spread patterns, and known local wildfire threat factors make it likely that a wildfire in this area would threaten the community. This includes stands with continuous surface/ crown fuel that will support regular torching/ candling, intermittent crown and/or continuous crown fires. Rates of spread would average 6 -10 meters/ minute. Fuel type spot potential is likely to impact values at a long distance (400 -1 000m).

Extreme (Red): The combination of the local fuel hazard, weather influences, topography, proximity to the community, fuel position in relation to fire spread patterns, and known local wildfire threat factors make it very likely that a wildfire in this area would threaten the community. Stands with continuous surface/ crown fuel and fuel characteristics that tend to support the development of intermittent or continuous crown fires. Rates of spread would average >10 meters/ minute. Fuel type spot potential is probable to impact values at a long distance (400 -1 000m or greater). These forest stands have the greater potential to produce extreme fire behaviour (long range spotting, fire whirls and other fire behaviour phenomena).

From Province Of British Columbia Community Wildfire Protection Plan Template 2017